

### Arthroscopic Capsular release Rehabilitation Protocol


#### JARED M. MAHYLIS, MD

Henry Ford Health System Detroit, MI

#### **Phase I**: (0 to 2 weeks after surgery)

Goals: Initiate Motion


- 1. Patients may remove dressing and shower 5 days after surgery. You may get the incision wet.
- 2. Sutures will be removed in clinic at your 1<sup>st</sup> follow up visit. These should be left in place.
- 3. Ice is recommended to reduce swelling and help with pain. You should continue this for the first week at a minimum. Apply the ice to the surgical site 3-4 times per day once the nerve block has worn off.
- 4. You may wear your sling when in a crowd, but stop wearing it once comfortable without it .
  - a. Initiate the following exercise program 3 times per day by the 1<sup>st</sup> postoperative day:
    - i. Immediate elbow, forearm, wrist, and hand range of motion out of sling. You may perform nonresistance, non-weight bearing use of the hand for activities such as typing on a computer or using a tablet computer.
    - ii. Start very gentle pendulum exercises.
    - iii. Begin active and active assisted: 1) forward elevation, 2) external rotation, 3) internal rotation, 4) extension and 5) cross body adduction. (see corresponding number for depiction of exercise.




## Phase II: (2 to 6 weeks after surgery)

Goals: Progress ROM & Strength

- 1. Continue Range of motion exercises
- 2. No lifting weights of more than 10-15 on your operative arm
- 3. Advance active and passive ROM in all planes to tolerance.
  - a. ER at the side and flexion to tolerance
  - b. Scapular plane elevation to 180



- c. IR and extension to tolerance
- Initiate gentle rotator cuff strengthening (see below)



# Phase III: (6 to 8 weeks after surgery)

Goals: Return to Full Function

- 1. Stop all lifting restrictions.
- 2. Progress on your rotator cuff and scapular stabilizer strengthening.
- 3. Initiate functional progression to activities.